*Journal of Biochemical and Biophysical Methods*, 24 (1992) 321–323

© 1992 Elsevier Science Publishers B.V. All rights reserved 0165-022X/92/$05.00

# Author Index


Alemany, M., see Latres, E. (24) 39
Altekar, W., see D'Souza, S.E. (24) 239
Ashizawa, K. and Cheng, S.-y.
A sensitive and rapid in situ immunoassay to quantitatively determine the cellular antigens in intact cultured cells (24) 297

Bertolini, J., see Schmitt, J.F. (24) 119
Boe, I.N., Lovrien, R.E. and Karl, D.W.
Stripping interfering sugars from samples using adapted bacteria (24) 309
Bøg-Hansen, T., see Kirkeby, S. (24) 225
Boschetti, E., see Santambien, P. (24) 285
Bridelli, M.G., Capelletti, R. and Vecli, A.
Sequential hydration-dehydration studies of lysozyme by the thermally stimulated depolarization currents (TSDC) technique (24) 135
Buettner, G.R. and Kiminyo, K.P.
Optimal EPR detection of weak nitroxide spin adduct and ascorbyl free radical signals (24) 147
Butko, P., Hapala, I., Nemecz, G. and Schroeder, F.
Sterol domains in phospholipid membranes: dehydroergosterol polarization measures molecular sterol transfer (24) 15

Cachia, C., see Flamion, P.-J. (24) 1
Capelletti, R., see Bridelli, M.G. (24) 135
Cebrian-Perez, J.A., see Pascual, M.L. (24) 275
Chang, G.-G., see Lee, H.-J. (24) 205
Cheng, S.-y., see Ashizawa, K. (24) 297
Chiou, S.-H., see Lee, H.-J. (24) 205
Chrambach, A., see Fawcett, J.S. (24) 181
Chrambach, A., see Wheeler, D. (24) 171
Cserháti, T. and Magyar, K.
Relationship between the hydrophobic and hydrophilic molecular parameters of some monoamine oxidase inhibitory drugs, determined by means of adsorptive and reversed-phase thin-layer chromatography (24) 249
Cummings, R.D., see Do, S.-I. (24) 153

Do, S.-I. and Cummings, R.D.
The hamster transferrin receptor contains Ser/Thr-linked oligosaccharides: use of a lectin-resistant CHO cell line to identify glycoproteins containing these linkages (24) 153
D'Souza, S.E., Altekar, W. and D'Souza, S.F.
A novel technique for the preparation of osmotically stabilized and permeabilized cells of extremely halophilic bacteria (24) 239
D'Souza, S.F., see D'Souza, S.E. (24) 239

Economou, C., see Schmitt, J.F. (24) 119

Fawcett, J.S., Wheeler, D. and Chrambach, A.
Transverse agarose pore gradient gel electrophoresis of DNA (24) 181
Fernández-López, J.-A., see Latres, E. (24) 39
Flamion, P.-J., Cachia, C. and Schreiber, J.-P.
Non-linear least-squares methods applied to the analysis of fluorescence energy transfer measurements (24) 1
Fountoulakis, M., Juranville, J.-F. and Manneberg, M.
Comparison of the Coomassie brilliant blue, bicinchoninic acid and Lowry quantitation assays, using non-glycosylated and glycosylated proteins (24) 265

Garbarsch, C., see Kirkeby, S. (24) 225
Garner, M.M., see Wheeler, D. (24) 171
Girot, P., see Santambien, P. (24) 285
Gosselin, S., Martin, B.M., Murray, G.J. and Viswanatha, T.
*Flavobacterium meningosepticum* peptide: *N*-glycosidase: influence of ionic strength on enzymatic activity (24) 71
Gu, H., Wilson, D. and Inselburg, J.
Recovery of DNA from agarose gels using a modified Elutrap™ (24) 45
Guthridge, M., see Schmitt, J.F. (24) 119

Hammerstedt, R.H., see Schweisguth, D.C. (24) 81
Hapala, I., see Butko, P. (24) 15

Hazum, E., Shisheva, A. and Shechter, Y.
Preparation and application of radioiodinated sulfhydryl reagents for the covalent labeling of SH-proteins present in minute quantities (24) 95
Hearn, M.T.W., see Schmitt, J.F. (24) 119
Hulak, I., see Santambien, P. (24) 285

Inouye, K., see Morimoto, K. (24) 107
Inselburg, J., see Gu, H. (24) 45

Juranville, J.-F., see Fountoulakis, M. (24) 265

Kachar, B., see Urrutia, R. (24) 63
Karl, D.W., see Boe, I.N. (24) 309
Katahira, M., see Nakamura, K.-i. (24) 195
Kiminyo, K.P., see Buettner, G.R. (24) 147
Kirkeby, S., Moe, D., Bøg-Hansen, T., Garbarsch, C. and Matthiessen, M.E.
Quantitative PAS assay of some carbohydrate compounds and detergents (24) 225

Latres, E., Fernández-López, J.-A., Remesar, X. and Alemany, M.
An enzymatic method for the estimation of L-leucine in rat blood (24) 39
Lee, H.-J., Chiou, S.-H. and Chang, G.-G.
Staining of argininosuccinate lyase activity in polyacrylamide gel (24) 205
Lee, V.S. and Tarassenko, L.
An optical method for the determination of platelet count in platelet samples contaminated with red blood cells (24) 215
Liras, A.
A rapid reliable method to use specific probes labelling polymerase chain reaction (PCR) products (24) 167
Lopez-Perez, M.J., see Pascual, M.L. (24) 275
Lovrien, R.E., see Boe, I.N. (24) 309

Magyar, K., see Cserháti, T. (24) 249
Manneberg, M., see Fountoulakis, M. (24) 265
Márquez-Méndez, M.
Purification of carboxypeptidase B by metal chelation affinity chromatography (24) 51
Martin, B.M., see Gosselin, S. (24) 71
Matthiessen, M.E., see Kirkeby, S. (24) 225
Moe, D., see Kirkeby, S. (24) 225
Morimoto, K. and Inouye, K.
Single-step purification of $F(ab')_2$ fragments of mouse monoclonal antibodies (immunoglobulins G1) by hydrophobic interaction high performance liquid chromatography using TSKgel Phenyl-5PW (24) 107
Muiño-Blanco, T., see Pascual, M.L. (24) 275
Murray, G.J., see Gosselin, S. (24) 71

Nakamura, K.-i., Okuya, Y., Katahira, M., Yoshida, S., Wada, S. and Okuno, M.
Analysis of tubulin isoforms by two-dimensional gel electrophoresis using SDS-polyacrylamide gel electrophoresis in the first dimension (24) 195
Nemecz, G., see Butko, P. (24) 15

Okuno, M., see Nakamura, K.-i. (24) 195
Okuya, Y., see Nakamura, K.-i. (24) 195
Orban, L., see Wheeler, D. (24) 171

Pascual, M.L., Muiño-Blanco, T., Cebrian-Perez, J.A. and Lopez-Perez, M.J.
Management of sedimentation in centrifugal counter-current distribution of sperm cells in an aqueous 2-phase system (24) 275

Remesar, X., see Latres, E. (24) 39

Santambien, P., Girot, P., Hulak, I. and Boschetti, E.
Immunochemical quantification of Procion Red HE-3B used as ligand in affinity chromatography (24) 285
Schmitt, J.F., Guthridge, M., Economou, C., Bertolini, J. and Hearn, M.T.W.
A new quantitative polymerase chain reaction-high performance ion exchange liquid chromatographic method for the detection of fibroblast growth factor-$\beta$ (FGF-$\beta$) gene amplification (24) 119
Schreiber, J.-P., see Flamion, P.-J. (24) 1
Schroeder, F., see Butko, P. (24) 15
Schweisguth, D.C. and Hammerstedt, R.H.
Evaluation of plasma membrane stability by detergent-induced rupture of osmotically swollen sperm (24) 81
Shechter, Y., see Hazum, E. (24) 95
Shisheva, A., see Hazum, E. (24) 95

Tarassenko, L., see Lee, V.S. (24) 215

Urrutia, R. and Kachar, B.
An improved method for the purification of kinesin from bovine adrenal medulla (24) 63

Vecli, A., see Bridelli, M.G. (24) 135
Viswanatha, T., see Gosselin, S. (24) 71

Wada, S., see Nakamura, K.-i. (24) 195
Wheeler, D., Orban, L., Garner, M.M. and Chrambach, A.
Computer-aided analysis of DNA curves on transverse gradient gels (24) 171
Wheeler, D., see Fawcett, J.S. (24) 181
Wilson, D., see Gu, H. (24) 45

Yoshida, S., see Nakamura, K.-i. (24) 195

*Journal of Biochemical and Biophysical Methods*, 24 (1992) 325–326

© 1992 Elsevier Science Publishers B.V. All rights reserved 0165-022X/92/$05.00

# Subject Index

Adapted bacteria, (24) 309
Adrenal medulla, (24) 63
Adsorption capacity, (24) 249
Affinity chromatography, (24) 285
Agarose, (24) 45
Agarose gradient, transverse, (24) 181
Aminoacyl-tRNA synthetase, (24) 39
Analytical HPLC, (24) 119
Argininosuccinate lyase, (24) 205
Ascorbyl radical, (24) 147
Axoneme, (24) 195

Bacterial stripping, (24) 309

Carbohydrate, (24) 225
Carboxypeptidase, (24) 51
Cell permeabilization, (24) 239
Cellular antigen, detection, (24) 297
Cholesterol, (24) 15
Colorimetric protein assay, (24) 265
Counter-current distribution, (24) 275
Covalent labeling, (24) 95
Cross-linking, (24) 239
Cryopreservation, (24) 81

δ-Crystallin, (24) 205
Deglycosylation, (24) 71
Dehydration kinetic, (24) 135
Dehydroergosterol, (24) 15
Detergent, (24) 81; (24) 225
Dextran, (24) 275
DNA, (24) 119; (24) 181
DNA cloning, (24) 167
DNA, conformation, (24) 171; (24) 181
DNA fragment, (24) 171
DNA purification, (24) 45
Dye leakage, (24) 285

EGF receptor, (24) 297
Electrodialysis, (24) 45
Electroelution, (24) 45
Electron paramagnetic resonance, (24) 147
Electrophoresis, agarose gradient, (24) 181
Electrophoresis, pore gradient, (24) 171
Electrophoresis, semi-automated, (24) 171; (24) 181
ELISA, (24) 285
Endocytosis, (24) 297
Enzyme staining, (24) 205
Epididymal maturation, (24) 81
Exchange, (24) 15
Extraction, (24) 45

$F(ab')_2$ fragment, (24) 107
Fatty-acid-binding-protein, (24) 15
Ferguson curve, computer traced, (24) 181
Ferguson curves, computer traced, (24) 171
Fibroblast growth factor, (24) 119
Fluorescence, (24) 15
Fluorescence energy transfer, (24) 1
Free radical, (24) 147

Gel electrophoresis, (24) 205
Glycerol, (24) 81
Glycoprotein, (24) 71

Halobacterium, (24) 239
*Helix pomatia* agglutinin, HPA, (24) 153
Hydration of proteins, (24) 135
Hydrophobic interaction, (24) 51
Hydrophobic interaction HPLC, (24) 107
Hypoosmotic, (24) 81

Immunoassay, in situ, (24) 297
Immunoglobulin G1, (24) 107
Ionic strength, (24) 71
Isoform, (24) 195

Kinesin, (24) 63

Least-squares, (24) 1
Lipophilicity, (24) 249
L-leucine, (24) 39
Lysis, (24) 81

Membrane, (24) 15; (24) 81
Metal chelate affinity chromatography, (24) 51
Monoamine oxidase inhibitor, (24) 249
Monoclonal antibody, (24) 107

Nitroxide radical, (24) 147

Organelle motility, (24) 63
Osmotic stabilization, (24) 239

Partitioning, (24) 275
Pepsin digestion, (24) 107
Peptide, (24) 1
Peptide: *N*-glycosidase, (24) 71
Percoll, (24) 275
Periodic acid-Schiff, PAS, (24) 225
Platelet count, (24) 215
Polarization, (24) 15
Poly(ethylene glycol), (24) 275
Polymerase chain reaction, (24) 119; (24) 167
Pore gradient, transverse, (24) 171
Probe labelling, (24) 167
Protein glycosylation, (24) 265
Protein purification, (24) 63
Pyruvate kinase $M_2$, (24) 297

Quantitation, (24) 119

Radioiodination, (24) 95
Recovery method, (24) 45
Resistance pulse spectrometry, (24) 81
tRNA, (24) 39

SH-protein, (24) 95
Single-step purification, (24) 107
Spectrophotometry, (24) 215
Spectroscopy, (24) 225
Spermatozoa, (24) 81; (24) 275
Spin trapping, (24) 147
Sterol-carrier protein, (24) 15
Sugar, (24) 309
Sulfhydryl reagent, (24) 95

*Tetrahymena*, (24) 195
Thermally stimulated depolarisation current, (24) 135
Thyroid hormone-binding protein, (24) 297
Transferrin receptor, TfR, (24) 153
Tryptophan, (24) 1
Tubulin, (24) 195
Two-dimensional gel electrophoresis, (24) 195
Tyrosine, (24) 1

Water sorption, (24) 135